

TESTfacts

An Educational Health Series From National Jewish Health®

SuPrep for Colonoscopy Liquid Diet

Start a liquid (no solid foods) diet from the moment of waking up on the day before the procedure.

Clear Liquid Diet

- Gatorade, fruit juice (no citrus), soda, popsicles without fruit or cream, gelatin without fruit, broth without fat, coffee, and tea
- No solid food
- No red, orange, or purple colored liquids
- No dairy

SuPrep Preparation Instructions:

- 1st dose of SuPrep to be taken at 6pm day of liquid diet. Mix 1 bottle from kit with water in the mixing cup supplied. Follow with 2 mixing cups of water.
- 2nd dose of SuPrep 6 hours prior to procedure. Mix second bottle from kit with water in the mixing cup supplied. Follow with 2 mixing cups of water.
- May have clear liquids up to 4 hours before the procedure.

Do not eat or drink (not even water) anything 4 hours prior to your procedure.

- Be sure to inform your doctors and nurses about any diabetic medications and/or blood thinners you take. This will be the last liquid you have.
- Take all medications for high blood pressure, heart conditions, or seizure disorder with water 4 hours prior to your procedure.
- Bring rescue inhalers, asthma inhalers, diabetic medications, glucose monitor, and CPAP or BiPap machine with you to the procedure.
- Please be sure to have a ride arranged and with you to take you home after the procedure. If you do not have a ride with you, your procedure may be canceled.

Questions? Please call (303)398-1355 option #4, if you have any questions or need to reschedule. ©National Jewish Health, PTE316, rev2/15

REGISTER NOW

to Receive FREE Health Tips, Research Findings, Ways to Support Our Mission, News & More in Your E-Mail Box

Visit njhealth.org/e-news for more information.

Follow us online!

facebook facebook.com/NJHealth



